

II. Synthetic Division

Can synthetic division be used to divide a polynomial by $x^2 - 5$? Explain.


NO, BECAUSE OF x^2

SYNTHETIC DIVISION ONLY WORKS WHEN
DIVIDING BY LINEAR TERMS.

Can synthetic division be used to divide a polynomial by $x + 4$? Explain.

YES, $x+4$ IS A LINEAR TERM

To divide $ax^3 + bx^2 + cx + d$ by $x - k$, use the following pattern:



III. The Remainder and Factor Theorems

To use the Remainder Theorem to evaluate a polynomial function $f(x)$ at $x = k$, you:

USE SYNTHETIC DIVISION TO DIVIDE $f(x)$ BY $x - k$. THE REMAINDER WILL BE $f(k)$

To use the Factor Theorem to show that $(x - k)$ is a factor of a polynomial function $f(x)$, you:

USE SYNTHETIC DIVISION ON $f(x)$ WITH THE FACTOR $x - k$. IF

THE REMAINDER IS 0, THEN $x - k$ IS A FACTOR OF $f(x)$.

List three facts about the remainder r , obtained in the synthetic division of $f(x)$ by $x - k$:

1) THE REMAINDER R GIVES THE VALUE OF f AT $x = k$. $r = f(k)$

2) IF $r = 0$, THEN $(x - k)$ IS A FACTOR OF $f(x)$.

3) IF $r = 0$, THEN $(k, 0)$ IS AN x -INTERCEPT OF THE GRAPH OF f .

What you should learn:

How to use synthetic division to divide polynomials by binomials of the form $(x - k)$

What you should learn:

How to use the Remainder and Factor Theorems

IV. The Rational Zero Test

Describe the purpose of the Rational Zero Test:

TO FIND POSSIBLE RATIONAL ZEROS OF
A POLYNOMIAL FUNCTION

What you should learn:

How to use the Rational Zero Test to determine possible rational zeros of polynomial functions

State the Rational Zero Test:

IF A POLYNOMIAL FUNCTION $f(x)$ HAS INTEGER COEFFICIENTS,
THEN EVERY RATIONAL ZERO OF f HAS THE FORM

$$\text{RATIONAL ZERO} = \frac{P}{Q}$$

\leftarrow FACTORS OF CONSTANT
 \leftarrow FACTORS OF LEADING COEFFICIENT

To use the Rational Zero Test, you:

- ① LIST ALL RATIONAL FACTORS OF p AND q
- ② TRIAL AND ERROR TO FIND ZEROS OF THE POLYNOMIAL

V. Other Tests for Zeros of Polynomials

State Descartes's Rule of Signs:

LET $f(x)$ BE A POLYNOMIAL WITH
REAL COEFFICIENTS:

- ① POSITIVE REAL ZEROS
 \leq TO NUMBER OF SIGN CHANGES (DECREASE BY 2)
- ② NEGATIVE REAL ZEROS
 \leq TO NUMBER OF SIGN CHANGES
IN $f(-x)$ (DECREASE BY 2)

What you should learn:

How to use Descartes's Rule of Signs and the Upper and Lower Bound Rules to find zeros of polynomials

Section 2.3 Examples – Real Zeros of Polynomial Functions

(1) Use long division to divide.

$$(x + 8 + 6x^3 + 10x^2) \div (2x^2 + 1)$$

$$\begin{array}{r}
 3x+5 \\
 2x^2+0x+1 \overline{) 6x^3+10x^2+x+8} \\
 \underline{-(6x^3+0x^2+3x)} \\
 10x^2-2x+8 \\
 \underline{-(10x^2+0x+5)} \\
 -2x+3
 \end{array}$$

$$\boxed{3x+5 + \frac{-2x+3}{2x^2+1}}$$

Labels: QUOTIENT (points to $3x+5$), REMAINDER (points to $-2x+3$), DIVISOR (points to $2x^2+1$)

(2) Use synthetic division to divide.

$$(3x^3 - 17x^2 + 15x - 25) \div (x - 5)$$

$$\begin{array}{r|rrrr}
 5 & 3 & -17 & 15 & -25 \\
 & \downarrow & 15 & -10 & 25 \\
 \hline
 & 3 & -2 & 5 & 0
 \end{array}$$

Labels: QUOTIENT COEFFICIENTS (points to 3, -2, 5), REMAINDER (points to 0)

$$\boxed{3x^2 - 2x + 5}$$

(3) Use the Remainder Theorem and Synthetic Division to evaluate the function at each given value.

$$g(x) = 2x^6 + 3x^4 - x^2 + 3$$

a) $g(2) = 175$

b) $g(1) = 7$

c) $g(3) = 1695$

d) $g(-1) = 7$

$$\begin{array}{r|rrrrrrr}
 2 & 2 & 0 & 3 & 0 & -1 & 0 & 3 \\
 & & 4 & 8 & 22 & 44 & 86 & 172 \\
 \hline
 & 2 & 4 & 11 & 22 & 43 & 86 & 175
 \end{array}$$

Label: REMAINDER (points to 175)

(4) Find ALL real zeros of the polynomial function.

$$f(x) = 4x^3 + 7x^2 - 11x - 18$$

$$\frac{P}{Q} = \pm \frac{1 \ 2 \ 3 \ 6 \ 9 \ 18}{1 \ 2 \ 4}$$

$$\begin{array}{r|rrrr}
 -2 & 4 & 7 & -11 & -18 \\
 & & -8 & 2 & 18 \\
 \hline
 & 4 & -1 & -9 & 0
 \end{array}$$

$$4x^2 - x - 9 = 0$$

$$x = \frac{1 \pm \sqrt{1+144}}{8}$$

$$\boxed{= \frac{1 \pm \sqrt{145}}{8}}$$